

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
12 February 2004 (12.02.2004)

PCT

(10) International Publication Number
WO 2004/012760 A1

- (51) International Patent Classification⁷: **A61K 38/30**,
A61P 19/08, 19/10
- (74) Agent: **BALDWIN SHELSTON WATERS**; P O Box
5999, Wellesley Street, 1000 Auckland (NZ).
- (21) International Application Number:
PCT/NZ2003/000168
- (22) International Filing Date: 31 July 2003 (31.07.2003)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
60/400,443 1 August 2002 (01.08.2002) US
- (71) Applicant (*for all designated States except US*):
AUCKLAND UNISERVICES LIMITED [NZ/NZ];
70 Symonds Street, 1000 AUCKLAND (NZ).
- (72) Inventors; and
- (75) Inventors/Applicants (*for US only*): **CORNISH, Jillian**
[NZ/NZ]; 23 Joseph Banks Terrace, Newmarket, Auckland
(NZ). **REID, Ian, Reginald** [NZ/NZ]; 7 Maybeck Road,
Mount Albert, Auckland (NZ). **COOPER, Garth, James,**
Smith [NZ/NZ]; 2 Crummer Road, Ponsonby, Auckland
(NZ). **BUCHANAN, Christina, Maree** [NZ/NZ]; 15 Cres-
cent Road West, Palm Beach, Waiheke, Auckland (NZ).
- (81) Designated States (*national*): AE, AG, AL, AM, AT (util-
ity model), AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA,
CH, CN, CO (utility model), CR, CU, CZ (utility model),
CZ, DE (utility model), DE, DK (utility model), DK, DM,
DZ, EC, EE (utility model), EE, ES, FI (utility model), FI,
GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE,
KG, KP, KR (utility model), KZ, LC, LK, LR, LS, LT, LU,
LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ,
OM, PG, PH, PL, PT (utility model), PT, RO, RU, SC, SD,
SE, SG, SK (utility model), SK, SL, SY, TJ, TM, TN, TR,
TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.
- (84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,
ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO,
SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM,
GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- Published:
— with international search report
- For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.*

WO 2004/012760 A1

(54) Title: PREPTIN METHODS OF USE

(57) Abstract: This invention features a method for treating a bone condition in a patient, e.g., a mammal, a human, a horse, a dog, or a cat. The method includes administering an effective amount of preptin, preptin analog, or a preptin agonist to the patient.

BEST AVAILABLE COPY

Preptin Methods of Use

BACKGROUND

Preptin is a 34-amino acid peptide corresponding to Asp⁶⁹-Leu¹⁰² of the proinsulin-like growth factor II. It is present in pancreatic islet beta cells and undergoes glucose-mediated co-secretion with insulin.

5 The two-phase secretion of insulin in response to glucose is well characterized. See, e.g., Grodsky *et al.* (1968) *Acta Diabetol. Lat.* 5: 140-161. The first phase results in a transient spike in insulin secretion, while the second phase results in a progressive increase in insulin release. Preptin has been found to enhance, but not initiate, insulin secretion. More specifically, infusion of preptin into the isolated, perfused rat pancreas exerts a significant
10 increase in the second phase of secretion, while removal of preptin from the pancreas by adding anti-preptin antibodies results in a decrease in both the first and the second phases of insulin secretion. See Buchanan *et al.* (2001) *Biochem. J.* 360: 431-439.

It also has been suggested that preptin elicits its effects by binding to a cell surface receptor (Buchanan *supra*).

15

SUMMARY

The present invention is based, in part, on the discovery that preptin can stimulate proliferation of osteoblasts, which are known to play a role in mediating or modulating bone growth.

20 In one aspect, this invention features a method for treating a bone condition in a patient, e.g., a mammal, a human, a horse, a dog, or a cat. The method includes administering an effective amount of preptin, preptin analog, or a preptin agonist to the patient.

25 The patient can be suffering from a disease associated with excessive resorption or breakdown of bone tissue. Examples of such diseases include, but are not limited to, osteoporosis, osteopenia, bone defects, and osteogenesis imperfecta. The patient can also be suffering from bone loss as a result of immobility, bone fractures, malignancy, primary hyperparathyroidism, endocrine disorders, autoimmune arthritis, or addictive drug use. The patient can also be undergoing a treatment (e.g., corticosteroid treatment, bone marrow

TJ504467/142

2

transplantation, or oophorectomy) known to result in bone loss. The term "bone condition" refers to any disease or symptom wherein mediation of osteoblast or osteoclast activity (or levels) is involved, and includes any of the diseases or situations described above.

As used herein, "preptin" is an isolated peptide of 34 amino acids in length, the
 5 sequence of which is described as in formula (I):

Asp Val Ser Thr R₁ R₂ R₃ Val Leu Pro Asp R₄ Phe Pro Arg Tyr Pro Val Gly Lys Phe
 Phe R₅ R₆ Asp Thr Trp R₇ Gln Ser R₈ R₉ Arg Leu formula (I)

Wherein:

R₁ is Ser or Pro
 10 R₂ is Gln or Pro
 R₃ is Ala or Thr
 R₄ is Asp or Asn
 R₅ is Gln or Lys
 R₆ is Tyr or Phe
 15 R₇ is Arg or Lys
 R₈ is Ala or Thr, and
 R₉ is Gly or Gln;
 or an analog thereof.

It includes mouse preptin, rat preptin, and human preptin, the sequences of which are shown
 20 below.

Mouse preptin: DVSTSQAVLPDDFPRYPVGKFFQYDTWRQSAGRL (SEQ ID NO: 1)

Rat preptin: DVSTSQAVLPDDFPRYPVGKFFKFDTWQRQSAGRL (SEQ ID NO: 2)

Human preptin: DVSTPPTVLPDNFPRYPVGKFFQYDTWKQSTQRL (SEQ ID NO: 3)

The amino acid sequence corresponds to Asp₆₉-Leu₁₀₂ of the proIGF-II E-peptide in each
 25 mammal.

Analogues of preptin include functional equivalents of preptin (e.g., functional
 equivalents of those of formula I). In terms of preptin itself, functional equivalents include
 all proteins which are immunologically cross-reactive with and have substantially the same
 function as preptin (e.g., any of SEQ ID NOs: 1-3). That equivalent may, for example, be a
 30 fragment of preptin containing from 6-33 amino acids (usually representing a C-terminal

TJ504467/142

3

truncation) and including a preptin active site or sites, a substitution, addition or deletion mutant of preptin, or a fusion of preptin or a fragment or a mutant with other amino acids.

“A preptin agonist” is a compound which (1) has a high affinity (e.g., a K_i of 10^{-7} - 10^{-9} M, a K_i of 10^{-8} - 10^{-9} M) for a preptin-binding receptor (as defined by the receptor binding assay described in Motulsky, H.J and Mahan, L.C. (1984). *Mol. Pharmacol.* 25: 1; and (2) promotes the proliferation of bone cells, e.g., osteoblasts.

In one embodiment, the methods described herein include administering to a patient an effective amount of preptin having the amino acid sequence of SEQ ID NO: 1, 2, or 3.

In another embodiment, the method includes administering to a patient an effective amount of a preptin agonist having a fragment (e.g., any sequence between 17 and 33 amino acids in length, inclusive, of SEQ ID NO: 1, 2, or 3) or the entirety of the amino acid sequence of SEQ ID NO: 1, 2, or 3. For example, a preptin agonist is a peptide being 17-110 amino acids in length, e.g., a peptide having less than 87 amino acids, or more than 20 (e.g., any integer between 21 and 110, inclusive) amino acids, and containing, in consecutive sequence, any part of SEQ ID NO: 1, 2, or 3. In another example, a preptin agonist is a peptide having 35-110 amino acids in length, e.g., a peptide having less than 87 amino acids, or more than 35 (e.g., any integer between 35 and 110, inclusive) amino acids, and containing the entirety of the amino acid sequence of SEQ ID NO: 1, 2, or 3.

In a further embodiment, the method includes administering to a patient an effective amount of a preptin agonist containing an amino acid sequence that is at least 60% (e.g., 70%, 80%, 90%, 95%, or 98%) identical to SEQ ID NO: 1, 2, or 3. The “percent identity” of two amino acid sequences can be determined using the algorithm of Karlin and Altschul (1990, *Proc. Natl. Acad. Sci. USA* 87: 2264-2268), modified as in Karlin and Altschul (1993, *Proc. Natl. Acad. Sci. USA* 90: 5873-5877). Such an algorithm is incorporated into the NBLAST and XBLAST programs (version 2.0) of Altschul *et al.* (1990) *J. Mol. Biol.* 215: 403-10. BLAST protein searches can be performed with the XBLAST program, score=50, wordlength=3 to obtain amino acid sequences homologous to the peptide molecules described herein. Where gaps exist between two sequences, Gapped BLAST can be utilized as described in Altschul *et al.* (1997) *Nucleic Acids Res.* 25(17): 3389-3402. When utilizing BLAST and Gapped BLAST programs, the default parameters of the respective programs (e.g., XBLAST and NBLAST) can be used.

TJ504467/142

4

In still another embodiment, the method includes administering to a patient an effective amount of a preptin agonist containing SEQ ID NO: 1, 2, or 3 with up to 14 (e.g., any integer between 1 and 14, inclusive) conservative amino acid substitutions. A “conservative amino acid substitution” is one in which an amino acid residue is replaced with another residue having a chemically similar side chain. Families of amino acid residues having similar side chains have been defined in the art. These families include amino acids with basic side chains (e.g., lysine, arginine, histidine), acidic side chains (e.g., aspartic acid, glutamic acid), uncharged polar side chains (e.g., glycine, asparagine, glutamine, serine, threonine, tyrosine, cysteine), nonpolar side chains (e.g., alanine, valine, leucine, isoleucine, proline, phenylalanine, methionine, tryptophan), beta-branched side chains (e.g., threonine, valine, isoleucine) and aromatic side chains (e.g., tyrosine, phenylalanine, tryptophan, histidine). Amino acid analogs (e.g., phosphorylated amino acids) are also contemplated in the present invention.

In another aspect, this invention features a method for increasing or maintaining bone density. The method includes administering to a subject (e.g., a mammal, a human, a horse, a dog, or a cat) in need thereof an effective amount of preptin, preptin analog, or a preptin agonist as described herein. As used herein, the subject may have a substantially normal bone density or the subject may be at risk of bone deterioration. Examples of these subjects include postmenopausal women, usually at age 50 and over, and men over 60 years of age.

In a further aspect, this invention features a method for stimulating osteoblast growth or modulating osteoblast apoptosis. The method includes administering to a subject in need thereof an effective amount of preptin, preptin analog, or a preptin agonist. The term “osteoblast” refers to bone-forming cells.

The method includes administering to the subject (including a subject identified as in need of such treatment, e.g., a subject in need of modulation of osteoblast activity) an effective amount of a compound described herein, or a composition described herein to produce such effect. Identifying a subject in need of such treatment can be in the judgment of a subject or a health care professional and can be subjective (e.g. opinion) or objective (e.g. measurable by a test or diagnostic method).

This invention also features an article of manufacture that includes a vessel containing preptin, preptin analog, or a preptin agonist; and instructions for use of preptin,

preptin analog, or a preptin agonist for treatment of a bone condition by administering an effective amount of preptin, preptin analog, or a preptin agonist to a patient.

Also within the scope of this invention is an article of manufacture. The article includes packaging material; and contained within the packaging material, preptin, preptin analog, or a preptin agonist. The packaging material comprises a label that indicates that preptin, preptin analog, or a preptin agonist can be used for treating a bone condition (e.g., osteoporosis, osteopenia, bone defects, or osteogenesis imperfecta) in a patient. In other aspects, the label includes dosage information.

Other features, objects, and advantages of the invention will be apparent from the description and drawings, and from the claims.

DESCRIPTION OF DRAWINGS

FIG. 1 depicts the effect of various concentrations of rat preptin or vehicle on cell number (FIG. 1A) and thymidine incorporation (FIG. 1B) in cultures of fetal rat osteoblasts. In this figure, * represents $p < 0.05$, ** represents $p < 0.01$, and *** represents $p < 0.001$.

FIG. 2 depicts the effect of various concentrations of human preptin or vehicle on cell number (FIG. 1A) and thymidine incorporation (FIG. 1B) in cultures of fetal rat osteoblasts. In this figure, * represents $p < 0.05$, ** represents $p < 0.01$, and *** represents $p < 0.001$.

FIG. 3 depicts the effects of two concentrations of rat preptin on thymidine incorporation in murine neonatal calvarial organ culture.

FIG. 4 depicts the effects of various concentrations of rat preptin on osteoclast development *in vitro*.

FIG. 5 depicts the effects of preptin on primary rat osteoblasts pre-treated with either MAP kinase inhibitors PD-98059 or U-0126 (FIG. 5A) or a G_i protein inhibitor, pertussis toxin (FIG. 5B).

FIG. 6 depicts the effects of rat preptin on apoptosis of serum-deprived primary rat osteoblasts.

DETAILED DESCRIPTION

This invention relates to use of preptin, preptin analog, or a preptin agonist for stimulating osteoblast growth or modulating osteoblast apoptosis. Preptin can be isolated

from pancreatic islet beta cells in a manner as described in Buchanan *et al.* (2001) *Biochem. J.* 360: 431-439. Preptin, as well as a preptin analog or preptin agonist, also can be prepared by a synthetic method. More specifically, synthesis of peptides (e.g., preptin) is well established in the art. See, e.g., Stewart, *et al.* (1984) *Solid Phase Peptide Synthesis* (2nd Ed.); and Chan (2000) "Fmoc Solid Phase Peptide Synthesis, A Practical Approach," Oxford University Press. The peptides may be synthesized using an automated peptide synthesizer (e.g., a Pioneer™ Peptide Synthesizer, Applied Biosystems, Foster City, CA). For example, a peptide is prepared on methylbenzylhydramine resin followed by hydrogen fluoride deprotection and cleavage from the resin. The synthesized peptide can be further purified by a method such as affinity column chromatography or high pressure liquid chromatography. Standard physicochemical characterization techniques are known in the art, including NMR (¹³C, ¹H, ¹⁹F, or ³¹P) and IR, which can provide confirmatory evidence of the identity and purity of the synthetic products. Amino acid analysis can also be used to confirm the amino acid composition of the peptide. Mass spectroscopy can be used to identify the molecular weight of synthetic products.

One aspect of this invention is a method for treating a bone condition with an effective amount of a preptin, preptin analog, or a preptin agonist. Another aspect of this invention is a method for increasing or maintaining bone density with a preptin, preptin analog, or a preptin agonist. The term "treating" is defined as the application or administration of a composition including a preptin, preptin analog, or a preptin agonist to a patient, who has, or is determined to have, a bone condition, a symptom of a bone condition, a disease or disorder secondary to a bone condition, or a predisposition toward a bone condition, with the purpose to cure, alleviate, relieve, remedy, or ameliorate the bone condition, the symptom of the bone condition, the disease or disorder secondary to the bone condition, or the predisposition toward the bone condition. "An effective amount" refers to an amount of preptin, preptin analog, or a preptin agonist that confers a therapeutic effect on the treated subject. The therapeutic effect may be objective (i.e., measurable by some test or marker) or subjective (i.e., subject gives an indication of or feels an effect). An effective amount of preptin, preptin analog, or a preptin agonist described above may range from about 1 Tg/Kg body weight to about 1000 Tg/Kg body weight. Effective doses will also vary depending on the route of administration, as well as the possibility of co-usage with other

agents for stimulating osteoblast growth or modulating osteoblast apoptosis, such as a bone anti-resorptive agent (e.g., calcitonin or bisphosphonate) or a bone anabolic agent (e.g., parathyroid hormone, parathyroid hormone related protein, cytokines, or growth hormone).

As used herein, preptin, preptin analog, and preptin agonists are defined to include
5 pharmaceutically acceptable derivatives (e.g., salts).

Pharmaceutically acceptable salts include those derived from pharmaceutically acceptable inorganic and organic acids and bases. Examples of suitable acid salts include acetate, adipate, alginate, aspartate, benzoate, benzenesulfonate, bisulfate, butyrate, citrate, camphorate, camphorsulfonate, cyclopentanepropionate, digluconate, dodecylsulfate,
10 ethanesulfonate, formate, fumarate, glucoheptanoate, glycerophosphate, glycolate, hemisulfate, heptanoate, hexanoate, hydrochloride, hydrobromide, hydroiodide, 2-hydroxyethanesulfonate, lactate, maleate, malonate, methanesulfonate, 2-naphthalenesulfonate, nicotinate, nitrate, palmoate, pectinate, persulfate, 3-phenylpropionate, phosphate, picrate, pivalate, propionate, salicylate, succinate, sulfate, tartrate, thiocyanate,
15 tosylate, trifluoroacetate, and undecanoate. Other acids, such as oxalic, while not in themselves pharmaceutically acceptable, may be employed in the preparation of salts useful as intermediates in obtaining the compounds of the invention and their pharmaceutically acceptable acid addition salts. Salts derived from appropriate bases include alkali metal (e.g., sodium), alkaline earth metal (e.g., magnesium), ammonium and N-(alkyl)₄⁺ salts. This
20 invention also envisions the quaternization of any basic nitrogen-containing groups of the compounds disclosed herein. Water or oil-soluble or dispersible products may be obtained by such quaternization.

Also within the scope of this invention is a pharmaceutical composition that contains an effective amount of preptin, preptin analog, or a preptin agonist, and a pharmaceutically
25 acceptable carrier.

The term "pharmaceutically acceptable carrier" refers to a carrier (adjuvant or vehicle) that may be administered to a patient, together with preptin, preptin analog, or a preptin agonist, and which does not destroy the pharmacological activity thereof and is nontoxic when administered in doses sufficient to deliver preptin, preptin analog, or a preptin
30 agonist.

Pharmaceutically acceptable carriers that may be used in the pharmaceutical compositions described above include, but are not limited to, ion exchangers, alumina, aluminum stearate, lecithin, self-emulsifying drug delivery systems (SEDDS) such as d- α -tocopherol polyethyleneglycol 1000 succinate, surfactants used in pharmaceutical dosage forms such as Tweens or other similar polymeric delivery matrices, serum proteins, such as human serum albumin, buffer substances such as phosphates, glycine, sorbic acid, potassium sorbate, partial glyceride mixtures of saturated vegetable fatty acids, water, salts or electrolytes, such as protamine sulfate, disodium hydrogen phosphate, potassium hydrogen phosphate, sodium chloride, zinc salts, colloidal silica, magnesium trisilicate, polyvinyl pyrrolidone, cellulose-based substances, polyethylene glycol, sodium carboxymethylcellulose, polyacrylates, waxes, polyethylene-polyoxypropylene-block polymers, polyethylene glycol and wool fat. Cyclodextrins such as α -, β -, and γ -cyclodextrin, or chemically modified derivatives such as hydroxyalkylcyclodextrins, including 2- and 3-hydroxypropyl- β -cyclodextrins, or other solubilized derivatives may also be advantageously used to enhance delivery of compounds of the formulae described herein. Oil solutions or suspensions may also contain a long-chain alcohol diluent or dispersant, or carboxymethyl cellulose or similar dispersing agents, which are commonly used in the formulation of pharmaceutically acceptable dosage forms such as emulsions and or suspensions.

To practice the method for treating a bone condition or the method for increasing or maintaining bone density, preptin, preptin analog, or a preptin agonist can be administered to a patient or a subject. The preptin, preptin analog, or the preptin agonist can, for example, be administered in a pharmaceutically acceptable carrier such as physiological saline, in combination with other drugs, and/or together with appropriate excipients. It also can, for example, be administered by injection, intravenously, intraarterially, subdermally, intraperitoneally, intramuscularly, or subcutaneously; or orally, buccally, nasally, transmucosally, topically, in an ophthalmic preparation, by inhalation, by intracranial injection or infusion techniques. The methods herein contemplate administration of an effective amount of compound or compound composition to achieve the desired or stated effect. Lower or higher doses than those described above may be required. Specific dosage and treatment regimens for any particular patient will depend upon a variety of factors,

including the activity of the specific compound employed, the age, body weight, general health status, sex, diet, time of administration, rate of excretion, drug combination, the severity and course of the disease, condition or symptoms, the patient's disposition to the disease, condition or symptoms, and the judgment of the treating physician.

5 A pharmaceutical composition may be orally administered in any orally acceptable dosage form including, but not limited to, capsules, tablets, emulsions and aqueous suspensions, dispersions and solutions. In the case of tablets for oral use, carriers that are commonly used include lactose and corn starch. Lubricating agents, such as magnesium stearate, are also typically added. For oral administration in a capsule form, useful diluents
10 include lactose and dried corn starch. When aqueous suspensions and/or emulsions are administered orally, the active ingredient may be suspended or dissolved in an oily phase is combined with emulsifying and/or suspending agents. If desired, certain sweetening and/or flavoring and/or coloring agents may be added.

 A sterile injectable composition (e.g., aqueous or oleaginous suspension) can be
15 formulated according to techniques known in the art using suitable dispersing or wetting agents (such as, for example, Tween 80) and suspending agents.

 Topical administration of a pharmaceutical composition is useful when the desired treatment involves areas or organs readily accessible by topical application. For application topically to the skin, the pharmaceutical composition should be formulated with a suitable
20 ointment containing the active components suspended or dissolved in a carrier. Carriers for topical administration of the compounds of this invention include, but are not limited to, mineral oil, liquid petroleum, white petroleum, propylene glycol, polyoxyethylene polyoxypropylene compound, emulsifying wax and water. Alternatively, the pharmaceutical composition can be formulated with a suitable lotion or cream containing the active
25 compound suspended or dissolved in a carrier with suitable emulsifying agents. Suitable carriers include, but are not limited to, mineral oil, sorbitan monostearate, polysorbate 60, cetyl esters wax, cetearyl alcohol, 2-octyldodecanol, benzyl alcohol and water. The pharmaceutical compositions of this invention may also be topically applied to the lower intestinal tract by rectal suppository formulation or in a suitable enema formulation.
30 Topically-applied transdermal patches are also included in this invention.

A pharmaceutical composition may be administered by nasal aerosol or inhalation. Such compositions are prepared according to techniques well-known in the art of pharmaceutical formulation and may be prepared as solutions in saline, employing benzyl alcohol or other suitable preservatives, absorption promoters to enhance bioavailability, fluorocarbons, and/or other solubilizing or dispersing agents known in the art.

Preptin agonists can be tested for their abilities to stimulate osteoblast growth or modulate osteoblast apoptosis by examining their activities in the *in vitro* assays described herein. See the specific examples below. *In vivo* screening can also be performed by following procedures well known in the art. See, e.g., Cornish *et al.* (1997) *Am J Physiol* 273: E1113-E1120; and Cornish *et al.* (2000) *Am J Physiol* 279: E730-E735.

All references cited herein, whether in print, electronic, computer readable storage media or other form, are expressly incorporated by reference in their entirety, including but not limited to, abstracts, articles, journals, publications, texts, treatises, internet web sites, databases, patents, and patent publications.

The invention will be further described in the following examples. It should be understood that these examples are for illustrative purposes only and are not to be construed as limiting this invention in any manner.

Example 1. Promoting proliferation of bone cells

Osteoblast-Like Cell Culture. Osteoblasts were isolated from 20 day fetal rat calvariae as previously described (Cornish *et al.* (1999) *American Journal of Physiology - Endocrinology & Metabolism* 277: E779-E783). Briefly, calvariae were excised and the frontal and parietal bones, free of suture and periosteal tissue, were collected. The calvariae were sequentially digested using collagenase and the cells from digests 3 and 4 were collected, pooled and washed. Cells were grown to confluence and then subcultured into 24 well plates. Cells were growth arrested in minimum essential medium (MEM)/0.1% bovine serum albumin for 24 h. Fresh media and experimental compounds were added for a further 24 h. Cells were pulsed with tritiated-thymidine two hours before the end of the experimental incubation. The effect of preptin on proliferation of fetal rat osteoblast-like cells was first assessed by the measurement of cell numbers. The effect of preptin on DNA synthesis in osteoblasts was then assessed by the measurement of [³H]-thymidine

incorporation into isolated fetal rat osteoblast-like cells. There were 6 wells in each group and each experiment was repeated 3 or 4 times.

As shown in FIGs 1 and 2, preptin, in a dose-dependent manner, stimulated the proliferation (cell number and DNA synthesis) of primary fetal rat osteoblasts and osteoblast-like cell lines at periphsiological concentrations ($>10^{-11}\text{M}$). In addition, thymidine
5 incorporation was stimulated in murine neonatal calvarial organ culture (FIG. 3), likely reflecting the proliferation of cells from the osteoblast lineage.

The effects of preptin on the development of osteoclasts was also examined. To assess osteoclast development, bone marrow is obtained from the long bones of normal mice,
10 aged 4-6 weeks and cultured. Non-adherent cells are removed and the cultures are grown in the presence of $1\alpha,25$ -dihydroxyvitamin D3 throughout the experiment. The cultures were maintained for 7 days and the number of tartrate-resistant acid phosphatase-positive multinucleated cells was assessed. An osteoclastogenesis assay was performed with murine bone marrow cultures in the presence of various concentrations of rat preptin. The number of
15 tartrate-resistant acid phosphatase (TRAP)-positive multinucleated cells was assessed (FIG. 4). Preptin did not affect osteoclast development in this assay. Similarly, preptin did not affect bone resorption in a murine neonatal calvarial organ culture system (data not shown).

20 Example 2. Inducing phosphorylation of p42/p44 MAP kinases in bone cells

Immunoblotting (Cell Signalling Methods). Primary rat osteoblasts prepared as described above were seeded in 6-well tissue culture plates at an initial density of 5×10^4 cells/mL in MEM 5% FCS, and grown to 80-90% confluence. After serum starvation overnight, cells were treated at room temperature with test substances in MEM 0.1% BSA.
25 In experiments designed to determine the effect of inhibitors of signal transduction on preptin-induced p42/44 MAP kinase phosphorylation, the cells were pre-treated with the inhibitor for 30 min prior to addition of test substances. The exception was pertussis toxin, which was added 18 h prior to test substances. After treatment for the indicated period of time, the treatment medium was aspirated, the cells were washed in ice-cold PBS and then
30 scraped in ice-cold HNTG lysis buffer (50 mM HEPES, pH 7.5, 150 mM NaCl, 1% Triton, 10% glycerol, 1.5 mM MgCl_2 , 1 mM EDTA) containing a cocktail of protease and

phosphatase inhibitors (1 mM PMSF, 1 µg/mL pepstatin, 10 µg/mL leupeptin, 10 µg/mL aprotinin, 1 mM sodium vanadate, 500 mM NaF). The lysates were briefly vortexed, clarified by centrifugation at 13,000 rpm at 4°C, then stored at -70°C until analyzed. Protein content of the cell lysates was measured using the DC protein assay (BioRad, Hercules, CA).
5 Equal amounts of whole cell lysate (30-50 µg) were subjected to 8% SDS-PAGE, transferred to nitrocellulose membranes and immunoblotted overnight at 4°C with an anti-phospho-p42/44 MAP kinase antibody (1:1000). As a control for protein loading, the same filters were stripped and re-probed with an antibody to total p42/44 MAP kinase (1:400).
Incubation with the HRP-conjugated secondary antibody was for 1 h at room temperature,
10 and the membranes were analyzed by ECL. *TUNEL assay (Apoptosis Method)*. Apoptosis in cultures of primary rat osteoblasts was assessed using the TUNEL method (DeadEnd™, Promega, Madison, WI) according to the manufacturer's instructions. Cells were seeded in 8-well chamber slides at 5×10^4 cells/mL in MEM containing 5% FCS. Twenty-four hours later, the medium was changed to MEM/0.1%BSA and the cells incubated overnight. The
15 following morning, test substances were added in fresh MEM/0.1% BSA for 18 h. At the end of the treatment period, cells were fixed in 2% paraformaldehyde for 15 min, then permeabilized with 1% Triton in PBS for 5 min. Thereafter, biotinylated nucleotides were added in the presence of terminal deoxynucleotidyl transferase for 1 h at 37°C and the reaction terminated with 2 x SSC. Endogenous peroxidases were blocked with 0.3% H₂O₂
20 for 5 min, streptavidin-HRP added for 30 min, and apoptotic nuclei colorized by addition of diaminobenzidine/H₂O₂ mixture. After counterstaining with hematoxylin, the number of apoptotic nuclei per microscopic field was counted and expressed as a proportion of that observed in the cells exposed to serum starvation throughout the entire treatment period. Each experiment was performed at least twice and involved assessment of at least 6 chambers
25 per treatment condition.

Preptin induced phosphorylation of p42/p44 MAP kinases in osteoblastic cells in a dose-dependent manner (10^{-8} - 10^{-10} M), as assessed by immunoblotting. The proliferative effects of preptin on primary osteoblasts were blocked when the cells were pre-treated with either of the MAP kinase kinase inhibitors PD-98059 or U-0126 (FIG. 5A), or a G_i protein inhibitor, pertussis toxin (FIG. 5B). The effect of preptin on primary osteoblast apoptosis
30 induced by serum deprivation was assessed. Apoptotic cells were detected by light

microscopy using a modified TUNEL assay. Preptin had anti-apoptotic effects, at 10^{-8} M with treatment/control ratio of 0.78 ± 0.08 (FIG. 6).

Example 3. Promoting bone growth *in vivo*

5 Several hormones that regulate nutritional status also impact on bone metabolism. Preptin, a 34-amino acid peptide hormone that increases glucose-mediated insulin secretion, has been recently isolated from the same secretory vesicles that contain insulin and amylin from the pancreatic β -cells. Preptin is anabolic to osteoblasts but, unlike amylin, does not regulate osteoclast activity.

10 Preptin not only stimulates osteoblast proliferation but also osteoblast differentiation at 10^{-8} M, significantly increasing the number of mineralized bone nodules in long-term osteoblast cultures. These effects are also seen *in vivo*, when preptin is injected locally over the hemicalvariae of sexually mature male mice. After five daily subcutaneous injections of 16.5 micrograms of preptin, there was a significant increase in bone area, and mineralizing surface.

Methods

20 Three groups of sexually mature male mice were given twice daily subcutaneous injections over the right hemicalvaria for 5 consecutive days. Two groups received one of 2 doses of preptin (0.825 μ g or 8.25 μ g per injection), and a further two group received vehicle. The animals were sacrificed 1 week following the last injection. Fluorochrome labels were injected subcutaneously at the base of the tail on days 1 (calcein), 5 (alizarin red) and 14 (calcein) to measure dynamic histomorphometric indices (such as, extent of mineralizing surface). Calvariae were excised, fixed in 10% neutral-buffered formalin, dehydrated and embedded in methylmethacrylate resin. Sections were cut, mounted on

25 gelatin-coated slides, and histomorphometric indices measured using image analysis.

Results

Local injections of high dose preptin significantly (by Student's *t* test) increased the bone area as well as the mineralizing surface compared to the control (see tables below).

Table 1. The effects of preptin on bone area *in vivo*

Bone Area			
	Control	Preptin (high)	Preptin (low)
Mean \pm Std Error	1.028 \pm 0.023	*1.084 \pm 0.017	1.083 \pm 0.027

*P = 0.013

Table 2. The effects of preptin on mineralizing surface of bone *in vivo*

% Mineralizing Surface/Total Bone Surface			
	Control	Preptin (high)	Preptin (low)
Mean \pm Std Error	1.048 \pm 0.013	*1.119 \pm 0.032	1.065 \pm 0.033

*P = 0.046

In conclusion, preptin, a peptide contained within proIGF-II, is anabolic to bone in *in vitro* and *in vivo* models. Since it is secreted from the pancreatic β -cell, it may act in concert with the other β -cell hormones, insulin and amylin, to stimulate bone formation in hyperinsulinemic states, such as obesity. Preptin may also contribute to the osteosclerotic phenotype observed in patients with chronic hepatitis C infection who have increased circulating levels of proIGF-II, which contains the preptin peptide (Khosla S, Ballard FJ, Conover CA. (2002) *J Clin Endocrinol Metab.* 87(8):3867-70). Thus, the anabolic effects of preptin seen in rodent models may influence bone density in humans.

OTHER EMBODIMENTS

All of the features disclosed in this specification may be combined in any combination. Thus, unless expressly stated otherwise, each feature disclosed is only an example of a generic series of equivalent or similar features.

It is to be understood that while the invention has been described in conjunction with the detailed description thereof, the foregoing description is intended to illustrate and not limit the scope of the invention, which is defined by the scope of the appended claims. Other aspects, advantages, and modifications are within the scope of the following claims.

CLAIMS:

1. A method for treating a bone condition, comprising administering to a patient in need thereof an effective amount of preptin, preptin analog, or a preptin agonist.
2. The method of claim 1, wherein the amino acid sequence of preptin is SEQ ID NO: 1, 2,
5 or 3.
3. The method of claim 1, wherein the preptin agonist comprises a fragment or the entirety of the amino acid sequence of SEQ ID NO: 1, 2, or 3.
- 10 4. The method of claim 3, wherein the fragment is amino acid residues 17-34 of SEQ ID NO: 1, 2, or 3.
5. The method of claim 1, wherein the preptin agonist comprises an amino acid sequence that is at least 60% identical to SEQ ID NO: 1, 2, or 3.
- 15 6. The method of claim 5, wherein the preptin agonist comprises an amino acid sequence that is at least 80% identical to SEQ ID NO: 1, 2, or 3.
7. The method of claim 5, wherein the preptin agonist comprises an amino acid sequence
20 that is at least 90% identical to SEQ ID NO: 1, 2, or 3.
8. The method of claim 5, wherein the preptin agonist comprises an amino acid sequence that is at least 95% identical to SEQ ID NO: 1, 2, or 3.
- 25 9. The method of claim 1, wherein the preptin agonist comprises SEQ ID NO: 1, 2, or 3 with up to 14 conservative amino acid substitutions.
10. The method of claim 9, wherein the preptin agonist comprises SEQ ID NO: 1, 2, or 3 with up to 10 conservative amino acid substitutions.

TJ504467/142

16

11. The method of claim 9, wherein the preptin agonist comprises SEQ ID NO: 1, 2, or 3 with up to 6 conservative amino acid substitutions.
12. The method of claim 9, wherein the preptin agonist comprises SEQ ID NO: 1, 2, or 3 with up to 2 conservative amino acid substitutions.
13. A method for increasing or maintaining bone density, comprising administering to a subject in need thereof an effective amount of preptin, preptin analog, or a preptin agonist.
14. The method of claim 13, wherein the amino acid sequence of preptin is SEQ ID NO: 1, 2, or 3.
15. The method of claim 13, wherein the preptin agonist comprises a fragment or the entirety of the amino acid sequence of SEQ ID NO: 1, 2, or 3.
16. The method of claim 15, wherein the fragment is amino acid residues 17-34 of SEQ ID NO: 1, 2, or 3.
17. The method of claim 13, wherein the preptin agonist comprises an amino acid sequence that is at least 60% identical to SEQ ID NO: 1, 2, or 3.
18. The method of claim 17, wherein the preptin agonist comprises an amino acid sequence that is at least 80% identical to SEQ ID NO: 1, 2, or 3.
19. The method of claim 17, wherein the preptin agonist comprises an amino acid sequence that is at least 90% identical to SEQ ID NO: 1, 2, or 3.
20. The method of claim 17, wherein the preptin agonist comprises an amino acid sequence that is at least 95% identical to SEQ ID NO: 1, 2, or 3.

TJ504467/142

17

21. The method of claim 13, wherein the preptin agonist comprises SEQ ID NO: 1, 2, or 3 with up to 14 conservative amino acid substitutions.
22. The method of claim 21, wherein the preptin agonist comprises SEQ ID NO: 1, 2, or 3 with up to 10 conservative amino acid substitutions.
23. The method of claim 21, wherein the preptin agonist comprises SEQ ID NO: 1, 2, or 3 with up to 6 conservative amino acid substitutions.
24. The method of claim 21, wherein the preptin agonist comprises SEQ ID NO: 1, 2, or 3 with up to 2 conservative amino acid substitutions.
25. A method for stimulating osteoblast growth or modulating osteoblast apoptosis, comprising administering to a subject in need thereof an effective amount of preptin, preptin analog, or a preptin agonist.
26. The method of claim 25, wherein the amino acid sequence of preptin is SEQ ID NO: 1, 2, or 3.
27. The method of claim 25, wherein the preptin agonist comprises a fragment or the entirety of the amino acid sequence of SEQ ID NO: 1, 2, or 3.
28. The method of claim 27, wherein the fragment is amino acid residues 17-34 of SEQ ID NO: 1, 2, or 3.
29. The method of claim 25, wherein the preptin agonist comprises an amino acid sequence that is at least 60% identical to SEQ ID NO: 1, 2, or 3.
30. The method of claim 29, wherein the preptin agonist comprises an amino acid sequence that is at least 80% identical to SEQ ID NO: 1, 2, or 3.

31. The method of claim 29, wherein the preptin agonist comprises an amino acid sequence that is at least 90% identical to SEQ ID NO: 1, 2, or 3.
32. The method of claim 29, wherein the preptin agonist comprises an amino acid sequence that is at least 95% identical to SEQ ID NO: 1, 2, or 3.
33. The method of claim 19, wherein the preptin agonist comprises SEQ ID NO: 1, 2, or 3 with up to 14 conservative amino acid substitutions.
34. The method of claim 33, wherein the preptin agonist comprises SEQ ID NO: 1, 2, or 3 with up to 10 conservative amino acid substitutions.
35. The method of claim 33, wherein the preptin agonist comprises SEQ ID NO: 1, 2, or 3 with up to 6 conservative amino acid substitutions.
36. The method of claim 33, wherein the preptin agonist comprises SEQ ID NO: 1, 2, or 3 with up to 2 conservative amino acid substitutions.
37. An article of manufacture comprising:
a vessel containing preptin, preptin analog, or a preptin agonist; and
instructions for use of preptin, preptin analog, or a preptin agonist for treatment of a bone condition comprising administering an effective amount of preptin, preptin analog, or a preptin agonist to a patient.
38. An article of manufacture comprising:
packaging material; and
contained within the packaging material, preptin, preptin analog, or a preptin agonist;
wherein the packaging material comprises a label that indicates that preptin, preptin analog, or a preptin agonist can be used for treating a bone condition in a patient.

39. Use of preptin, a preptin analog, or a preptin agonist in the manufacture of a medicament for treating a bone condition.
40. Use of preptin, a preptin analog, or a preptin agonist in the manufacture of a medicament
5 for increasing or maintaining bone density.
41. Use of preptin, a preptin analog, or a preptin agonist in the manufacture of a medicament for stimulating osteoblast growth or modulating osteoblast apoptosis.
- 10 42. Use according to any of claims 39 to 41, wherein the amino acid sequence of preptin is SEQ ID NO: 1, 2, or 3.
43. Use according to any one of claims 39 to 41, wherein the preptin agonist comprises a fragment or the entirety of the amino acid sequence of SEQ ID NO: 1, 2, or 3.
15
44. Use according to claim 43, wherein the fragment is amino acid residues 17-34 of SEQ ID NO: 1, 2, or 3.
45. Use according to any one of claims 39 to 41, wherein the preptin agonist comprises an
20 amino acid sequence that is at least 60% identical to SEQ ID NO: 1, 2, or 3.
46. Use according to claim 45, wherein the preptin agonist comprises an amino acid sequence that is at least 80% identical to SEQ ID NO: 1, 2, or 3.
- 25 47. Use according to claim 45, wherein the preptin agonist comprises an amino acid sequence that is at least 90% identical to SEQ ID NO: 1, 2, or 3.
48. Use according to claim 45, wherein the preptin agonist comprises an amino acid sequence that is at least 95% identical to SEQ ID NO: 1, 2, or 3.

49. Use according to any one of claims 39 to 41, wherein the preptin agonist comprises SEQ ID NO: 1, 2, or 3 with up to 14 conservative amino acid substitutions.
50. Use according to claim 49, wherein the preptin agonist comprises SEQ ID NO: 1, 2, or 3
5 with up to 10 conservative amino acid substitutions.
51. Use according to claim 49, wherein the preptin agonist comprises SEQ ID NO: 1, 2, or 3
with up to 6 conservative amino acid substitutions.
- 10 52. Use according to claim 49, wherein the preptin agonist comprises SEQ ID NO: 1, 2, or 3
with up to 2 conservative amino acid substitutions.




FIG. 1A

FIG. 1B




FIG. 2A




FIG. 2B

3/6




FIG. 3

4/6




FIG. 4

5/6




FIG. 5B

FIG. 5A

6/6




FIG. 6

INTERNATIONAL SEARCH REPORT

International application No.
PCT/NZ03/00168

A. CLASSIFICATION OF SUBJECT MATTER		
Int. Cl. ⁷ : A61K 38/30; A61P 19/08, 19/10		
According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED		
Minimum documentation searched (classification system followed by classification symbols)		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) Derwent, DWPI: preptin		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P, X	REID, I.R., "Relationships among Body Mass, Its Components, and Bone", BONE, (November 2002), Vol. 31, No. 5, pages 547-555 Entire document	37, 38
X	BUCHANAN, C.M. et al., "Preptin derived from proinsulin-like growth factor II (pro IGF-II) is secreted from pancreatic islet β -cells and enhances insulin secretion" BIOCHEM. J., (2001), 360, pages 431-439 Entire document	37, 38
X	WO 00/78805 A1 (COOPER, G.J.S. et al.) 28 December 2000 Entire document	37, 38
<input type="checkbox"/> Further documents are listed in the continuation of Box C <input checked="" type="checkbox"/> See patent family annex		
<p>* Special categories of cited documents:</p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier application or patent but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art</p> <p>"&" document member of the same patent family</p>		
Date of the actual completion of the international search 29 October 2003		Date of mailing of the international search report - 7 NOV 2003
Name and mailing address of the ISA/AU AUSTRALIAN PATENT OFFICE PO BOX 200, WODEN ACT 2606, AUSTRALIA E-mail address: pct@pastralia.gov.au Facsimile No. (02) 6285 3929		Authorized officer NICOLE HOWARD Telephone No : (02) 6283 2245

INTERNATIONAL SEARCH REPORT

International application No.

PCT/NZ03/00168

This Annex lists the known "A" publication level patent family members relating to the patent documents cited in the above-mentioned international search report. The Australian Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

Patent Document Cited in Search Report				Patent Family Member			
WO	0078805	AU	57178/00	CA	2375207	EP	1185558
		US	2003050434	US	2003166561		
END OF ANNEX							

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.